

# MASS FORMULA FOR SUPERSINGULAR ABELIAN SURFACES

CHIA-FU YU AND JENG-DAW YU

ABSTRACT. We show a mass formula for arbitrary supersingular abelian surfaces in characteristic  $p$ .

## 1. INTRODUCTION

In [1] Chai studied prime-to- $p$  Hecke correspondences on Siegel moduli spaces in characteristic  $p$  and proved a deep geometric result about ordinary  $\ell$ -adic Hecke orbits for any prime  $\ell \neq p$ . Recently Chai and Oort gave a complete answer to what this  $\ell$ -adic Hecke orbit can be; see [2]. In this paper we study the arithmetic aspect of supersingular  $\ell$ -adic Hecke orbits in the Siegel moduli spaces, the extreme situation opposite to the ordinary case. In the case of genus  $g = 2$ , we give a complete answer to the size of supersingular Hecke orbits.

Let  $p$  be a rational prime number and  $g \geq 1$  be a positive integer. Let  $N \geq 3$  be a prime-to- $p$  positive integer. Choose a primitive  $N$ th root of unity  $\zeta_N \in \overline{\mathbb{Q}} \subset \mathbb{C}$  and fix an embedding  $\overline{\mathbb{Q}} \hookrightarrow \overline{\mathbb{Q}_p}$ . Let  $\mathcal{A}_{g,1,N}$  denote the moduli space over  $\overline{\mathbb{F}_p}$  of  $g$ -dimensional principally polarized abelian varieties with a symplectic level- $N$  structure with respect to  $\zeta_N$ . Let  $k$  be an algebraically closed field of characteristic  $p$ . For each point  $x = \underline{A}_0 = (A_0, \lambda_0, \eta_0)$  in  $\mathcal{A}_{g,1,N}(k)$  and a prime number  $\ell \neq p$ , the  $\ell$ -adic Hecke orbit  $\mathcal{H}_\ell(x)$  is defined to be the countable subset of  $\mathcal{A}_{g,1,N}(k)$  that consists of points  $\underline{A}$  such that there is an  $\ell$ -quasi-isogeny from  $A$  to  $A_0$  that preserves the polarizations (see §2 for definitions). It is proved in Chai [1, Proposition 1] that the  $\ell$ -adic Hecke orbit  $\mathcal{H}_\ell(x)$  is finite if and only if  $x$  is supersingular. Recall that an abelian variety  $A$  over  $k$  is *supersingular* if it is isogenous to a product of supersingular elliptic curves;  $A$  is *superspecial* if it is isomorphic to a product of supersingular elliptic curves. A natural question is whether it is possible to calculate the size of a supersingular Hecke orbit. The answer is affirmative, provided that we know its underlying  $p$ -divisible group structure explicitly, through the calculation of geometric mass formulas (see Section 2). This is the task of this paper where we examine the  $p$ -divisible group structure of some non-superspecial abelian varieties.

Let  $x = (A_0, \lambda_0)$  be a  $g$ -dimensional supersingular principally polarized abelian varieties over  $k$ . Let  $\Lambda_x$  denote the set of isomorphism classes of  $g$ -dimensional supersingular principally polarized abelian varieties  $(A, \lambda)$  over  $k$  such that there exists an isomorphism  $(A, \lambda)[p^\infty] \simeq (A_0, \lambda_0)[p^\infty]$  of the attached quasi-polarized  $p$ -divisible groups; it is a finite set (see [7, Theorem 2.1 and Proposition 2.2]). Define the mass  $\text{Mass}(\Lambda_x)$  of  $\Lambda_x$  as

$$(1.1) \quad \text{Mass}(\Lambda_x) := \sum_{(A, \lambda) \in \Lambda_x} \frac{1}{|\text{Aut}(A, \lambda)|}.$$

---

*Date:* February 2, 2009.

The main result of this paper is computing the geometric mass  $\text{Mass}(\Lambda_x)$  for arbitrary  $x$  when  $g = 2$ .

Let  $\Lambda_{2,p}^*$  be the set of isomorphism classes of polarized superspecial abelian surfaces  $(A, \lambda)$  with polarization degree  $\deg \lambda = p^2$  over  $\overline{\mathbb{F}}_p$  such that  $\ker \lambda \simeq \alpha_p \times \alpha_p$  (see §3.1). For each member  $(A_1, \lambda_1)$  in  $\Lambda_{2,p}^*$ , the space of degree- $p$  isogenies  $\varphi : (A_1, \lambda_1) \rightarrow (A, \lambda)$  with  $\varphi^* \lambda = \lambda_1$  over  $k$  is a projective line  $\mathbf{P}^1$  over  $k$ . Write  $\mathbf{P}_{A_1}^1$  to indicate the space of  $p$ -isogenies arising from  $A_1$ . This family is studied in Moret-Bailly [6], and also in Katsura-Oort [5]. One defines an  $\mathbb{F}_{p^2}$ -structure on  $\mathbf{P}^1$  using the  $W(\mathbb{F}_{p^2})$ -structure of  $M_1$  defined by  $F^2 = -p$ , where  $M_1$  is the covariant Dieudonné module of  $A_1$  and  $F$  is the absolute Frobenius. For any supersingular principally polarized abelian surface  $(A, \lambda)$  there exist an  $(A_1, \lambda_1)$  in  $\Lambda_{2,p}^*$  and a degree- $p$  isogeny  $\varphi : (A_1, \lambda_1) \rightarrow (A, \lambda)$  with  $\varphi^* \lambda = \lambda_1$ . The choice of  $(A_1, \lambda_1)$  and  $\varphi$  may not be unique. However, the degree  $[\mathbb{F}_{p^2}(\xi) : \mathbb{F}_{p^2}]$  of the point  $\xi \in \mathbf{P}_{A_1}^1(k)$  that corresponds to  $\varphi$  is well-defined.

In this paper we prove

**Theorem 1.1.** *Let  $x = (A, \lambda)$  be a supersingular principally polarized abelian surface over  $k$ . Suppose that  $(A, \lambda)$  is represented by a pair  $(\underline{A}_1, \xi)$ , where  $\underline{A}_1 \in \Lambda_{2,p}^*$  and  $\xi \in \mathbf{P}_{A_1}^1(k)$ . Then*

$$\text{Mass}(\Lambda_x) = \frac{L_p}{5760},$$

where

$$L_p = \begin{cases} (p-1)(p^2+1) & \text{if } \mathbb{F}_{p^2}(\xi) = \mathbb{F}_{p^2}, \\ (p^2-1)(p^4-p^2) & \text{if } [\mathbb{F}_{p^2}(\xi) : \mathbb{F}_{p^2}] = 2, \\ (p^2-1)|\text{PSL}_2(\mathbb{F}_{p^2})| & \text{otherwise.} \end{cases}$$

Theorem 1.1 calculates the cardinality of  $\ell$ -adic Hecke orbits  $\mathcal{H}_\ell(x)$ , as one has (Corollary 2.3)

$$|\mathcal{H}_\ell(x)| = |\text{Sp}_{2g}(\mathbb{Z}/N\mathbb{Z})| \cdot \text{Mass}(\Lambda_x).$$

We mention that the function field analogue of Theorem 1.1 where supersingular abelian surfaces are replaced by supersingular Drinfeld modules is established in [10].

This paper is organized as follows. In Section 2 we describe the relationship between supersingular  $\ell$ -adic Hecke orbits and mass formulas. We develop the mass formula for the orbits of certain superspecial abelian varieties. In Section 3 we compute the endomorphism ring of any supersingular abelian surface. The proof of the main theorem is given in the last section.

## 2. HECKE ORBITS AND MASS FORMULAS

Let  $g, p, N, \ell, \mathcal{A}_{g,1,N}, k$  be as in the previous section. We work with a slightly bigger moduli space in which the objects are not necessarily equipped with principal polarizations. It is indeed more convenient to work in this setting. Let  $\mathcal{A}_{g,p^*,N} = \cup_{m \geq 1} \mathcal{A}_{g,p^m,N}$  be the moduli space over  $\overline{\mathbb{F}}_p$  of  $g$ -dimensional abelian varieties together with a  $p$ -power degree polarization and a symplectic level- $N$  structure with respect to  $\zeta_N$ . Write  $\mathcal{A}_{g,p^*}$  for the moduli stack over  $\overline{\mathbb{F}}_p$  that parametrizes  $g$ -dimensional  $p$ -power degree polarized abelian varieties. For any point  $x = \underline{A}_0 = (A_0, \lambda_0, \eta_0)$  in  $\mathcal{A}_{g,p^*,N}(k)$ , the  $\ell$ -adic Hecke orbit  $\mathcal{H}_\ell(x)$  is defined to be the countable subset of  $\mathcal{A}_{g,p^*,N}(k)$  that consists of points  $\underline{A}$  such that there

is an  $\ell$ -quasi-isogeny from  $A$  to  $A_0$  that preserves the polarizations. An  $\ell$ -quasi-isogeny from  $A$  to  $A_0$  is an element  $\varphi \in \text{Hom}(A, A_0) \otimes \mathbb{Q}$  such that  $\ell^m \varphi$ , for some integer  $m \geq 0$ , is an isogeny of  $\ell$ -power degree.

**2.1. Group theoretical interpretation.** Assume that  $x$  is supersingular. Let  $G_x$  be the automorphism group scheme over  $\mathbb{Z}$  associated to  $\underline{A}_0$ ; for any commutative ring  $R$ , the group of its  $R$ -valued points is defined by

$$G_x(R) = \{h \in (\text{End}_k(A_0) \otimes R)^\times \mid h'h = 1\},$$

where  $h \mapsto h'$  is the Rosati involution induced by  $\lambda_0$ . Let  $\Lambda_{x,N} \subset \mathcal{A}_{g,p^*,N}(k)$  be the subset consisting of objects  $(A, \lambda, \eta)$  such that there is an isomorphism  $\epsilon_p : (A, \lambda)[p^\infty] \simeq (A_0, \lambda_0)[p^\infty]$  of quasi-polarized  $p$ -divisible groups. Since  $\ell$ -quasi-isogenies do not change the associated  $p$ -divisible group structure, we have the inclusion  $\mathcal{H}_\ell(x) \subset \Lambda_{x,N}$ .

**Proposition 2.1.** *Notations and assumptions as above.*

- (1) *There is a natural isomorphism  $\Lambda_{x,N} \simeq G_x(\mathbb{Q}) \backslash G_x(\mathbb{A}_f) / K_N$  of pointed sets, where  $K_N$  is the stabilizer of  $\eta_0$  in  $G_x(\mathbb{Z})$ .*
- (2) *One has  $\mathcal{H}_\ell(x) = \Lambda_{x,N}$ .*

PROOF. (1) This is a special case of [7, Theorem 2.1 and Proposition 2.2]. We sketch the proof for the reader's convenience. Let  $\underline{A}$  be an element in  $\Lambda_{x,N}$ . As  $A$  is supersingular, there is a quasi-isogeny  $\varphi : A_0 \rightarrow A$  such that  $\varphi^* \lambda = \lambda_0$ . For each prime  $q$  (including  $p$  and  $\ell$ ), choose an isomorphism  $\epsilon_q : \underline{A}_0[q^\infty] \simeq \underline{A}[q^\infty]$  of  $q$ -divisible groups compatible with polarizations and level structures. There is an element  $\phi_q \in G_x(\mathbb{Q}_q)$  such that  $\varphi \phi_q = \epsilon_q$  for all  $q$ . The map  $\underline{A} \mapsto [(\phi_q)]$  gives a well-defined map from  $\Lambda_{x,N}$  to  $G_x(\mathbb{Q}) \backslash G_x(\mathbb{A}_f) / K_N$ . It is not hard to show that this is a bijection.

- (2) The inclusion  $\mathcal{H}_\ell(x) \subset \Lambda_{x,N}$  under the isomorphism in (1) is given by

$$[G_x(\mathbb{Q}) \cap G_x(\hat{\mathbb{Z}}^{(\ell)})] \backslash [G_x(\mathbb{Q}_\ell) \times G_x(\hat{\mathbb{Z}}^{(\ell)})] / K_N \subset G_x(\mathbb{Q}) \backslash G_x(\mathbb{A}_f) / K_N.$$

Since the group  $G_x$  is semi-simple and simply-connected, the strong approximation shows that  $G_x(\mathbb{Q}) \subset G_x(\mathbb{A}_f^{(\ell)})$  is dense. The equality then follows immediately. ■

**Corollary 2.2.** *Let  $\underline{A}_i = (A_i, \lambda_i, \eta_i)$ ,  $i = 1, 2$ , be two supersingular points in  $\mathcal{A}_{g,p^*,N}(k)$ . Suppose that there is an isomorphism of the associated quasi-polarized  $p$ -divisible groups. Then for any prime  $\ell \nmid pN$  there is an  $\ell$ -quasi-isogeny  $\varphi : A_1 \rightarrow A_2$  which preserves the polarizations and level structures.*

PROOF. This follows from the strong approximation property for  $G_x$  that any element  $\phi$  in the double space  $G_x(\mathbb{Q}) \backslash G_x(\mathbb{A}_f) / K_N$  can be represented by an element in  $G_x(\mathbb{Q}_\ell) \times K_N^{(\ell)}$ , where  $K_N^{(\ell)} \subset G_x(\hat{\mathbb{Z}}^{(\ell)})$  is the prime-to- $\ell$  component of  $K_N$ . ■

Recall that we denote by  $\Lambda_x$  the set of isomorphism classes of  $g$ -dimensional supersingular  $p$ -power degree polarized abelian varieties  $(A, \lambda)$  over  $k$  such that there is an isomorphism  $(A, \lambda)[p^\infty] \simeq (A_0, \lambda_0)[p^\infty]$ , and define the mass  $\text{Mass}(\Lambda_x)$  of  $\Lambda_x$  as

$$\text{Mass}(\Lambda_x) := \sum_{(A, \lambda) \in \Lambda_x} \frac{1}{|\text{Aut}(A, \lambda)|}.$$

Similarly, we define

$$\text{Mass}(\Lambda_{x,N}) := \sum_{(A,\lambda,\eta) \in \Lambda_{x,N}} \frac{1}{|\text{Aut}(A, \lambda, \eta)|}.$$

**Corollary 2.3.** *One has  $|\mathcal{H}_\ell(x)| = |\text{Sp}_{2g}(\mathbb{Z}/N\mathbb{Z})| \cdot \text{Mass}(\Lambda_x)$ .*

PROOF. This follows from

$$|\mathcal{H}_\ell(x)| = |\Lambda_{x,N}| = \text{Mass}(\Lambda_{x,N}) = |G_x(\mathbb{Z}/N\mathbb{Z})| \cdot \text{Mass}(\Lambda_x)$$

and  $|G_x(\mathbb{Z}/N\mathbb{Z})| = |\text{Sp}_{2g}(\mathbb{Z}/N\mathbb{Z})|$ . ■

**2.2. Relative indices.** Write  $G'$  for the automorphism group scheme associated to a principally polarized superspecial point  $x_0$ . The group  $G'_\mathbb{Q}$  is unique up to isomorphism. This is an inner form of  $\text{Sp}_{2g}$  which is “twisted at  $p$  and  $\infty$ ” (cf. §3.1 below). For any supersingular point  $x \in \mathcal{A}_{g,p^*}(k)$ , we can regard  $G_x(\mathbb{Z}_p)$  as an open compact subgroup of  $G'(\mathbb{Q}_p)$  through a choice of a quasi-isogeny of polarized abelian varieties between  $x_0$  and  $x$ . Another choice of quasi-isogeny gives rise to a subgroup which differs from the previous one by the conjugation of an element in  $G'(\mathbb{Q}_p)$ . For any two open compact subgroups  $U_1, U_2$  of  $G'(\mathbb{Q}_p)$ , we put

$$\mu(U_1/U_2) := [U_1 : U_1 \cap U_2][U_2 : U_1 \cap U_2]^{-1}.$$

**Proposition 2.4.** *Let  $x_1, x_2$  be two supersingular points in  $\mathcal{A}_{g,p^*}(k)$ . Then one has*

$$\text{Mass}(\Lambda_{x_2}) = \text{Mass}(\Lambda_{x_1}) \cdot \mu(G_{x_1}(\mathbb{Z}_p)/G_{x_2}(\mathbb{Z}_p)).$$

PROOF. See Theorem 2.7 of [7]. ■

**2.3. The superspecial case.** Let  $\Lambda_g$  denote the set of isomorphism classes of  $g$ -dimensional principally polarized superspecial abelian varieties over  $\overline{\mathbb{F}}_p$ . When  $g = 2D > 0$  is even, we denote by  $\Lambda_{g,p^D}^*$  the set of isomorphism classes of  $g$ -dimensional polarized superspecial abelian varieties  $(A, \lambda)$  of degree  $p^{2D}$  over  $\overline{\mathbb{F}}_p$  satisfying  $\ker \lambda = A[F]$ , where  $F : A \rightarrow A^{(p)}$  is the relative Frobenius morphism on  $A$ . Write

$$M_g := \sum_{(A,\lambda) \in \Lambda_g} \frac{1}{|\text{Aut}(A, \lambda)|}, \quad M_g^* := \sum_{(A,\lambda) \in \Lambda_{g,p^D}^*} \frac{1}{|\text{Aut}(A, \lambda)|}$$

for the mass attached to the finite sets  $\Lambda_g$  and  $\Lambda_{g,p^D}^*$ , respectively.

**Theorem 2.5.** *Notations as above.*

(1) *For any positive integer  $g$ , one has*

$$M_g = \frac{(-1)^{g(g+1)/2}}{2^g} \left\{ \prod_{k=1}^g \zeta(1-2k) \right\} \cdot \prod_{k=1}^g \{(p^k + (-1)^k)\},$$

where  $\zeta(s)$  is the Riemann zeta function.

(2) *For any positive even integer  $g = 2D$ , one has*

$$M_g^* = \frac{(-1)^{g(g+1)/2}}{2^g} \left\{ \prod_{k=1}^g \zeta(1-2k) \right\} \cdot \prod_{k=1}^D (p^{4k-2} - 1).$$

PROOF. (1) This is due to Ekedahl and Hashimoto-Ibukiyama (see [3, p.159] and [4, Proposition 9], also cf. [8, Section 3]).

(2) See Theorem 6.6 of [8]. ■

**Corollary 2.6.** *One has*

$$M_2 = \frac{(p-1)(p^2+1)}{5760}, \quad \text{and} \quad M_2^* = \frac{(p^2-1)}{5760}.$$

PROOF. This follows from Theorem 2.5 and the basic fact  $\zeta(-1) = \frac{-1}{12}$  and  $\zeta(-3) = \frac{1}{120}$ . This is also obtained in Katsura-Oort [5, Theorem 5.1 and Theorem 5.2] by a method different from above. ■

*Remark 2.7.* Proposition 2.1 is generalized to the moduli spaces of PEL-type in [9], with modification due to the failure of the Hasse principle.

### 3. ENDOMORPHISM RINGS

In this section we treat the endomorphism rings of supersingular abelian surfaces.

**3.1. Basic setting.** For any abelian variety  $A$  over  $k$ , the *a-number*  $a(A)$  of  $A$  is defined by

$$a(A) := \dim_k \operatorname{Hom}(\alpha_p, A).$$

Here  $\alpha_p$  is the kernel of the Frobenius morphism  $F : \mathbb{G}_a \rightarrow \mathbb{G}_a$  on the additive group. Denote by  $\mathcal{DM}$  the category of Dieudonné modules over  $k$ . If  $M$  is the (covariant) Dieudonné module of  $A$ , then

$$a(A) = a(M) := \dim_k M/(F, V)M.$$

Let  $B_{p,\infty}$  denote the quaternion algebra over  $\mathbb{Q}$  which is ramified exactly at  $\{p, \infty\}$ . Let  $D$  be the division quaternion algebra over  $\mathbb{Q}_p$  and  $O_D$  be the maximal order. Let  $W = W(k)$  be the ring of Witt vectors over  $k$ ,  $B(k) := \operatorname{Frac}(W(k))$  the fraction field, and  $\sigma$  the Frobenius map on  $W(k)$ . We also write  $\mathbb{Q}_{p^2}$  and  $\mathbb{Z}_{p^2}$  for  $B(\mathbb{F}_{p^2})$  and  $W(\mathbb{F}_{p^2})$ , respectively.

Let  $A$  be an abelian variety (over any field). The endomorphism ring  $\operatorname{End}(A)$  is an order of the semi-simple algebra  $\operatorname{End}(A) \otimes \mathbb{Q}$ . Determining  $\operatorname{End}(A)$  is equivalent to determining the semi-simple algebra  $\operatorname{End}(A) \otimes \mathbb{Q}$  and all local orders  $\operatorname{End}(A) \otimes \mathbb{Z}_\ell$ . Suppose that  $A$  is a supersingular abelian variety over  $k$ . We know that

- $\operatorname{End}(A) \otimes \mathbb{Q} = M_g(B_{p,\infty})$ , and
- $\operatorname{End}(A) \otimes \mathbb{Z}_\ell = M_{2g}(\mathbb{Z}_\ell)$  for all primes  $\ell \neq p$ .

Therefore, it is sufficient to determine the local endomorphism ring  $\operatorname{End}(A) \otimes \mathbb{Z}_p = \operatorname{End}_{\mathcal{DM}}(M)$ , which is an order of the simple algebra  $M_g(D)$ .

**3.2. The surface case.** Let  $A$  be a supersingular abelian surface over  $k$ . There is a superspecial abelian surface  $A_1$  and an isogeny  $\varphi : A_1 \rightarrow A$  of degree  $p$ . Let  $M_1$  and  $M$  be the covariant Dieudonné modules of  $A_1$  and  $A$ , respectively. One regards  $M_1$  as a submodule of  $M$  through the injective map  $\varphi_*$ . Let  $N$  be the Dieudonné submodule in  $M_1 \otimes \mathbb{Q}_p$  such that  $VN = M_1$ . If  $a(M) = 1$ , then  $M_1 = (F, V)M$  and hence it is determined by  $M$ . If  $a(M) = 2$ , or equivalently  $M$  is superspecial, then there are  $p^2 + 1$  superspecial submodules  $M_1 \subset M$  such that  $\dim_k M/M_1 = 1$ .

Now we fix a rank 4 superspecial Dieudonné module  $N$  (and hence fix  $M_1$ ) and consider the space  $\mathcal{X}$  of Dieudonné submodules  $M$  with  $M_1 \subset M \subset N$  and

$\dim_k N/M = 1$ . It is clear that  $\mathcal{X}$  is isomorphic to the projective line  $\mathbf{P}^1$  over  $k$ . Let  $\tilde{N} \subset N$  be the  $W(\mathbb{F}_{p^2})$ -submodule defined by  $F^2 = -p$ . This gives an  $\mathbb{F}_{p^2}$ -structure on  $\mathbf{P}^1$ . It is easy to show the following

**Lemma 3.1.** *Let  $\xi \in \mathbf{P}^1(k)$  be the point corresponding to a Dieudonné module  $M$  in  $\mathcal{X}$ . Then  $M$  is superspecial if and only if  $\xi \in \mathbf{P}^1(\mathbb{F}_{p^2})$ .*

Choose a  $W$ -basis  $e_1, e_2, e_3, e_4$  for  $N$  such that

$$Fe_1 = e_2, \quad Fe_2 = -pe_1, \quad Fe_3 = e_4, \quad Fe_4 = -pe_3.$$

Note that this is a  $W(\mathbb{F}_{p^2})$ -basis for  $\tilde{N}$ . Write  $\xi = [a : b] \in \mathbf{P}^1(k)$ . The corresponding Dieudonné module  $M$  is given by

$$M = \text{Span} \langle pe_1, pe_3, e_2, e_4, v \rangle,$$

where  $v = a'e_1 + b'e_3$  and  $a', b' \in W$  are any liftings of  $a, b$  respectively.

**Case (i):**  $\xi \in \mathbf{P}^1(\mathbb{F}_{p^2})$ . In this case  $M$  is superspecial. We have  $\text{End}_{\mathcal{DM}}(M) = M_2(O_D)$ .

Assume that  $\xi \notin \mathbf{P}^1(\mathbb{F}_{p^2})$ . In this case  $a(M) = 1$ . If  $\phi \in \text{End}_{\mathcal{DM}}(M)$ , then  $\phi \in \text{End}_{\mathcal{DM}}(N)$ . Therefore,

$$\text{End}_{\mathcal{DM}}(M) = \{ \phi \in \text{End}_{\mathcal{DM}}(N) ; \phi(M) \subset M \}.$$

We have  $\text{End}_{\mathcal{DM}}(N) = \text{End}_{\mathcal{DM}}(\tilde{N}) = M_2(O_D)$ . The induced map

$$(3.1) \quad \pi : \text{End}_{\mathcal{DM}}(\tilde{N}) \rightarrow \text{End}_{\mathcal{DM}}(\tilde{N}/V\tilde{N})$$

is surjective. Put

$$V_0 := \tilde{N}/V\tilde{N} = \mathbb{F}_{p^2}e_1 \oplus \mathbb{F}_{p^2}e_3 \quad \text{and} \quad B_0 := \text{End}_{\mathbb{F}_{p^2}}(V_0).$$

We have

$$\text{End}_{\mathcal{DM}}(\tilde{N}/V\tilde{N}) = \text{End}_{\mathbb{F}_{p^2}}(V_0) = M_2(\mathbb{F}_{p^2}).$$

Put

$$B'_0 := \{ T \in B_0 ; T(v) \in k \cdot v \},$$

where  $v = ae_1 + be_3 \in V_0 \otimes_{\mathbb{F}_{p^2}} k$ . Therefore,  $\text{End}_{\mathcal{DM}}(M) = \pi^{-1}(B'_0)$ . Since  $\xi \notin \mathbf{P}^1(\mathbb{F}_{p^2})$ ,  $a \neq 0$ . We write  $\xi = [1 : b]$ ,  $v = e_1 + be_3$ , and we have  $\mathbb{F}_{p^2}(\xi) = \mathbb{F}_{p^2}(b)$ .

Write  $T = \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} \in B_0$ , where  $a_{ij} \in \mathbb{F}_{p^2}$ . From  $T(v) \in kv$ , we get the condition

$$(3.2) \quad a_{12}b^2 + (a_{11} - a_{22})b - a_{21} = 0.$$

**Case (ii):**  $\mathbb{F}_{p^2}(\xi)/\mathbb{F}_{p^2}$  is quadratic. Write  $\xi = [1 : b]$ . Suppose  $b$  satisfies  $b^2 = \alpha b + \beta$ , where  $\alpha, \beta \in \mathbb{F}_{p^2}$ . Plugging this in (3.2), we get

$$a_{11} - a_{12} + a_{12}\alpha = 0 \quad \text{and} \quad a_{12}\beta = a_{21}.$$

This shows

$$(3.3) \quad B'_0 = \left\{ t_1 I + t_2 \begin{pmatrix} 0 & 1 \\ \beta & \alpha \end{pmatrix} ; t_1, t_2 \in \mathbb{F}_{p^2} \right\} \simeq \mathbb{F}_{p^2}(\xi),$$

where  $X^2 - \alpha X - \beta$  is the minimal polynomial of  $b$ .

**Case (iii):**  $\xi \notin \mathbf{P}^1(\mathbb{F}_{p^2})$  and  $\mathbb{F}_{p^2}(\xi)/\mathbb{F}_{p^2}$  is not quadratic. In this case  $a_{12} = a_{21} = 0$  and  $a_{11} = a_{22}$ . We have

$$B'_0 = \left\{ \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix} ; a \in \mathbb{F}_{p^2} \right\}.$$

We conclude

**Proposition 3.2.** *Let  $A$  be a supersingular surface over  $k$  and  $M$  be the associated covariant Dieudonné module. Suppose that  $A$  is represented by a pair  $(A_1, \xi)$ , where  $A_1$  is a superspecial abelian surface and  $\xi \in \mathbf{P}_{A_1}^1(k)$ . Let  $\pi : M_2(O_D) \rightarrow M_2(\mathbb{F}_{p^2})$  be the natural projection.*

- (1) *If  $\mathbb{F}_{p^2}(\xi) = \mathbb{F}_{p^2}$ , then  $\text{End}_{\mathcal{DM}}(M) = M_2(O_D)$ .*
- (2) *If  $[\mathbb{F}_{p^2}(\xi) : \mathbb{F}_{p^2}] = 2$ , then*

$$\text{End}_{\mathcal{DM}}(M) \simeq \{ \phi \in M_2(O_D) ; \pi(\phi) \in B'_0 \},$$

where  $B'_0 \subset M_2(\mathbb{F}_{p^2})$  is a subalgebra isomorphic to  $\mathbb{F}_{p^2}(\xi)$ .

- (3) *If it is not in the case (1) or (2), then*

$$\text{End}_{\mathcal{DM}}(M) \simeq \left\{ \phi \in M_2(O_D) ; \pi(\phi) = \begin{pmatrix} a & 0 \\ 0 & a \end{pmatrix}, a \in \mathbb{F}_{p^2} \right\}.$$

#### 4. PROOF OF THEOREM 1.1

**4.1. The automorphism groups.** Let  $x = (A, \lambda)$  be a supersingular principally polarized abelian surfaces over  $k$ . Let  $x_1 = (A_1, \lambda_1)$  be an element in  $\Lambda_{2,p}^*$  such that there is a degree- $p$  isogeny  $\varphi : (A_1, \lambda_1) \rightarrow (A, \lambda)$  of polarized abelian varieties. Write  $\xi = [a : b] \in \mathbf{P}^1(k)$  the point corresponding to the isogeny  $\varphi$ . We choose an  $\mathbb{F}_{p^2}$ -structure on  $\mathbf{P}^1$  as in §3.2. Let  $(M_1, \langle, \rangle) \subset (M, \langle, \rangle)$  be the covariant Dieudonné modules associated to  $\varphi : (A_1, \lambda_1) \rightarrow (A, \lambda)$ . Let  $N$  be the submodule in  $M_1 \otimes_{\mathbb{Q}_p}$  such that  $VN = M_1$ , and put  $\langle, \rangle_N = p\langle, \rangle$ . One has an isomorphism  $(N, \langle, \rangle_N) \simeq (M_1, \langle, \rangle)$  of quasi-polarized Dieudonné modules. Put

$$\begin{aligned} U_x &:= G_x(\mathbb{Z}_p) = \text{Aut}_{\mathcal{DM}}(M, \langle, \rangle), \\ U_{x_1} &:= G_{x_1}(\mathbb{Z}_p) = \text{Aut}_{\mathcal{DM}}(M_1, \langle, \rangle) = \text{Aut}_{\mathcal{DM}}(N, \langle, \rangle_N). \end{aligned}$$

Choose a  $W$ -basis  $e_1, e_2, e_3, e_4$  for  $N$  such that

$$\begin{aligned} Fe_1 &= e_2, & Fe_2 &= -pe_1, & Fe_3 &= e_4, & Fe_4 &= -pe_3, \\ \langle e_1, e_3 \rangle_N &= -\langle e_3, e_1 \rangle_N = 1, & \langle e_2, e_4 \rangle_N &= -\langle e_4, e_2 \rangle_N = p, \end{aligned}$$

and  $\langle e_i, e_j \rangle = 0$  for all remaining  $i, j$ . The Dieudonné module  $M$  is given by

$$M = \text{Span} \langle pe_1, pe_3, e_2, e_4, v \rangle,$$

where  $v = a'e_1 + b'e_3$  and  $a', b' \in W$  are any liftings of  $a, b$  respectively.

**Case (i):**  $\xi \in \mathbf{P}^1(\mathbb{F}_{p^2})$ . In this case  $A$  is superspecial. One has  $\Lambda_x = \Lambda_2$  and, by Corollary 2.6,

$$\text{Mass}(\Lambda_x) = \frac{(p-1)(p^2+1)}{5760}.$$

In the remaining of this section, we treat the case  $\xi \notin \mathbf{P}^1(\mathbb{F}_{p^2})$ . One has

$$U_x = \{ \phi \in U_{x_1} ; \phi(M) = M \},$$

and, by Proposition 2.4 and Corollary 2.6,

$$(4.1) \quad \text{Mass}(\Lambda_x) = \text{Mass}(\Lambda_{x_1}) \cdot \mu(U_{x_1}/U_x) = \frac{p^2-1}{5760} [U_{x_1} : U_x].$$

Recall that  $V_0 = \tilde{N}/V\tilde{N}$ , which is equipped with the non-degenerate alternating pairing  $\langle \cdot, \cdot \rangle : V_0 \times V_0 \rightarrow \mathbb{F}_{p^2}$  induced from  $\langle \cdot, \cdot \rangle_N$ . The map (3.1) induces a group homomorphism

$$\pi : U_{x_1} \rightarrow \text{Aut}(V_0, \langle \cdot, \cdot \rangle) = \text{SL}_2(\mathbb{F}_{p^2}).$$

**Proposition 4.1.** *The map  $\pi$  above is surjective.*

The proof is given in Subsection 4.2.

**Lemma 4.2.** *One has  $\ker \pi \subset U_x$ .*

PROOF. Let  $\phi \in \ker \pi$ . Write  $\phi(e_1) = e_1 + f_1, \phi(e_3) = e_3 + f_3$ , where  $f_1, f_3 \in VN$ . Since  $M$  is generated by  $VN$  and  $v$ , it suffices to check  $\phi(v) = v + a'f_1 + b'f_3 \in M$ ; this is clear. ■

**Case (ii):**  $[\mathbb{F}_{p^2}(\xi) : \mathbb{F}_{p^2}] = 2$ . By Proposition 3.2 and Lemma 4.2, we have  $\pi : U_{x_1}/U_x \simeq \text{SL}_2(\mathbb{F}_{p^2})/\mathbb{F}_{p^2}(\xi)_1^\times$ . This shows

$$[U_{x_1} : U_x] = (p^4 - p^2).$$

**Case (iii):**  $[\mathbb{F}_{p^2}(\xi) : \mathbb{F}_{p^2}] \geq 3$ . By Proposition 3.2 and Lemma 4.2, we have  $\pi : U_{x_1}/U_x \simeq \text{SL}_2(\mathbb{F}_{p^2})/\{\pm 1\}$ . This shows

$$[U_{x_1} : U_x] = |\text{PSL}_2(\mathbb{F}_{p^2})|.$$

From Cases (i)-(iii) above and equation (4.1), Theorem 1.1 is proved.

#### 4.2. Proof of Proposition 4.1. Write

$$O_D = W(\mathbb{F}_{p^2})[\Pi], \quad \Pi^2 = -p, \quad \Pi a = a^\sigma \Pi, \quad \forall a \in W(\mathbb{F}_{p^2}).$$

The canonical involution is given by  $(a + b\Pi)^* = a^\sigma - b\Pi$ . With the basis  $1, \Pi$ , we have the embedding

$$O_D \subset M_2(W(\mathbb{F}_{p^2})), \quad a + b\Pi = \begin{pmatrix} a & -pb^\sigma \\ b & a^\sigma \end{pmatrix}.$$

Note that this embedding is compatible with the canonical involutions. With respect to the basis  $e_1, e_2, e_3, e_4$ , an element  $\phi \in \text{End}_{\mathcal{DM}}(N)$  can be written as

$$T = (T_{ij}) \in M_2(O_D) \subset M_4(W(\mathbb{F}_{p^2})), \quad T_{ij} = a_{ij} + b_{ij}\Pi = \begin{pmatrix} a_{ij} & -pb_{ij}^\sigma \\ b_{ij} & a_{ij}^\sigma \end{pmatrix}.$$

Since  $\phi$  preserves the pairing  $\langle \cdot, \cdot \rangle_N$ , we get the condition in  $M_4(\mathbb{Q}_{p^2})$ :

$$(4.2) \quad T^t \begin{pmatrix} & J \\ -J & \end{pmatrix} T = \begin{pmatrix} & J \\ -J & \end{pmatrix}, \quad J = \begin{pmatrix} 1 & \\ & p \end{pmatrix}.$$

Note that

$$w_0 T_{ji}^* w_0^{-1} = T_{ji}^t, \quad w_0 = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(\mathbb{Z}_{p^2}).$$

The condition (4.2) becomes

$$(4.3) \quad \begin{pmatrix} w_0 & \\ & w_0 \end{pmatrix} T^* \begin{pmatrix} w_0^{-1} & \\ & w_0^{-1} \end{pmatrix} \begin{pmatrix} & J \\ -J & \end{pmatrix} T = \begin{pmatrix} & J \\ -J & \end{pmatrix}.$$



Since

$$\begin{pmatrix} w_0^{-1} & \\ & w_0^{-1} \end{pmatrix} \begin{pmatrix} & J \\ -J & \end{pmatrix} = \begin{pmatrix} & -\Pi \\ \Pi & \end{pmatrix} = \Pi \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(O_D),$$

we have

**Lemma 4.3.** *The group  $U_{x_1}$  is the group of  $O_D$ -linear automorphisms on the standard  $O_D$ -lattice  $O_D \oplus O_D$  which preserve that quaternion hermitian form  $\begin{pmatrix} 0 & -\Pi \\ \Pi & 0 \end{pmatrix}$ .*

We also write (4.3) as

$$(4.4) \quad \Pi^{-1} T^* \Pi w T = w, \quad w = \begin{pmatrix} & -1 \\ 1 & \end{pmatrix} \in M_2(O_D).$$

Notation. For an element  $T \in M_m(D)$  and  $n \in \mathbb{Z}$ , write  $T^{(n)} = \Pi^n T \Pi^{-n}$ . In particular, if  $T = (T_{ij}) \in M_m(\mathbb{Q}_{p^2}) \subset M_m(D)$ , then  $T^{(n)} = (T_{ij}^{\sigma^n})$ . If  $T \in M_m(O_D)$ , denote by  $\bar{T} \in M_m(\mathbb{F}_{p^2})$  the reduction mod  $\Pi$ .

Suppose we are given  $\bar{\phi} \in \text{SL}_2(\mathbb{F}_{p^2})$ , we must find an element  $T \in M_2(O_D)$  satisfying (4.4). We show that there is a sequence of elements  $T_n \in M_2(O_D)$  for  $n \geq 0$  satisfying the conditions

$$(4.5) \quad (T_n^*)^{(1)} w T_n \equiv w \pmod{\Pi^{n+1}}, \quad T_{n+1} \equiv T_n \pmod{\Pi^{n+1}}, \quad \text{and} \quad \bar{T}_0 = \bar{\phi}.$$

Suppose there is already an element  $T_n \in M_2(O_D)$  for some  $n \geq 0$  that satisfies

$$(T_n^*)^{(1)} w T_n \equiv w \pmod{\Pi^{n+1}}.$$

Put  $T_{n+1} := T_n + B_n \Pi^{n+1}$ , where  $B_n \in M_2(O_D)$ , and put  $X_n := (T_n^*)^{(1)} w T_n$ . Suppose  $X_n \equiv w + C_n \Pi^{n+1} \pmod{\Pi^{n+2}}$ . One computes that

$$\begin{aligned} X_{n+1} &\equiv T_n^{*(1)} w T_n + T_n^{*(1)} w B_n \Pi^{n+1} + (\Pi^{n+1})^* B_n^{*(1)} w T_n \pmod{\Pi^{n+2}} \\ &\equiv w + C_n \Pi^{n+1} + T_n^{*(1)} w B_n \Pi^{n+1} + (-1)^{n+1} B_n^{*(n)} w T_n^{(n+1)} \Pi^{n+1} \pmod{\Pi^{n+2}}. \end{aligned}$$

Therefore, we require an element  $B_n \in M_2(O_D)$  satisfying

$$\bar{C}_n + \bar{T}_n^t w \bar{B}_n + (-1)^{n+1} \bar{B}_n^{t(n+1)} w \bar{T}_n^{(n+1)} = 0.$$

Put  $\bar{Y}_n := \bar{T}_n^t w \bar{B}_n$ . As  $\bar{Y}_n^t = -\bar{B}_n^t w \bar{T}_n$ , we need to solve the equation

$$\bar{C}_n + \bar{Y}_n + (-1)^n \bar{Y}_n^{t(n+1)} = 0,$$

or equivalently the equation

$$\begin{cases} \bar{C}_n + \bar{Y}_n + \bar{Y}_n^{t(1)} = 0, & \text{if } n \text{ is even,} \\ \bar{C}_n + \bar{Y}_n - \bar{Y}_n^t = 0, & \text{if } n \text{ is odd.} \end{cases}$$

It is easy to compute that  $X_n^* = -X_n^{(1)}$ . From this it follows that

$$(-1)^{n+1} C_n^{*(n+1)} \Pi^{n+1} \equiv -C_n^{(1)} \Pi^{n+1} \pmod{\Pi^{n+2}},$$

or simply  $(-1)^n \bar{C}_n^{t(n)} = \bar{C}_n^{(1)}$ . This gives the condition

$$\begin{cases} \bar{C}_n^t = \bar{C}_n^{(1)}, & \text{if } n \text{ is even,} \\ -\bar{C}_n^t = \bar{C}_n, & \text{if } n \text{ is odd.} \end{cases}$$

By the following lemma, we prove the existence of  $\{T_n\}$  satisfying (4.5). Therefore, Proposition 4.1 is proved.

**Lemma 4.4.** *Let  $C$  be an element in the matrix algebra  $M_m(\mathbb{F}_{p^2})$ .*

- (1) *If  $C^t = C^{(1)}$ , then there is an element  $Y \in M_m(\mathbb{F}_{p^2})$  such that  $C + Y + Y^{t(1)} = 0$ .*
- (2) *If  $-C^t = C$ , then there is an element  $Y \in M_m(\mathbb{F}_{p^2})$  such that  $C + Y - Y^t = 0$ .*

PROOF. The proof is elementary and hence omitted. ■

*Remark 4.5.* Theorem 1.1 also provides another way to look at the supersingular locus  $S_2$  of the Siegel threefold. We used to divide it into two parts: superspecial locus and non-superspecial locus. Consider the mass function

$$M : S_2 \rightarrow \mathbb{Q}, \quad x \mapsto \text{Mass}(\Lambda_x).$$

Then the function  $M$  divides the supersingular locus  $S_2$  into 3 locally closed subsets that refine the previous one. More generally, we can consider the same function  $M$  on the supersingular locus  $S_g$  of the Siegel modular variety of genus  $g$ . The situation definitely becomes much more complicated. However, it is worth knowing whether the following question has the affirmative answer.

**(Question):** Is the map  $M : S_g \rightarrow \mathbb{Q}$  a constructible function?

*Acknowledgments.* The research of C.-F. Yu was partially supported by the grants NSC 97-2115-M-001-015-MY3 and AS-98-CDA-M01. The research of J.-D. Yu was partially supported by the grant NSC 97-2115-M-002-018.

#### REFERENCES

- [1] C.-L. Chai, Every ordinary symplectic isogeny class in positive characteristic is dense in the moduli. *Invent. Math.* **121** (1995), 439–479.
- [2] C.-L. Chai, Hecke orbits on Siegel modular varieties. *Geometric methods in algebra and number theory*, 71–107, *Progr. Math.*, **235**, Birkhauser Boston, 2005.
- [3] T. Ekedahl, On supersingular curves and supersingular abelian varieties. *Math. Scand.* **60** (1987), 151–178.
- [4] K. Hashimoto and T. Ibukiyama, On class numbers of positive definite binary quaternion hermitian forms, *J. Fac. Sci. Univ. Tokyo* **27** (1980), 549–601.
- [5] T. Katsura and F. Oort, Families of supersingular abelian surfaces, *Compositio Math.* **62** (1987), 107–167.
- [6] L. Moret-Bailly, Familles de courbes et de variétés abéliennes sur  $\mathbf{P}^1$ . Sém. sur les pinceaux de courbes de genre au moins deux (ed. L. Szpiro). *Astérisques* **86** (1981), 109–140.
- [7] C.-F. Yu, On the mass formula of supersingular abelian varieties with real multiplications. *J. Australian Math. Soc.* **78** (2005), 373–392.
- [8] C.-F. Yu, The supersingular loci and mass formulas on Siegel modular varieties. *Doc. Math.* **11** (2006), 449–468.
- [9] C.-F. Yu, Simple mass formulas on Shimura varieties of PEL-type. To appear in *Forum Math.*
- [10] C.-F. Yu and J. Yu, Mass formula for supersingular Drinfeld modules. *C. R. Acad. Sci. Paris Sér. I Math.* **338** (2004) 905–908.

(C.-F. Yu) INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, 128 ACADEMIA RD. SEC. 2, NANKANG, TAIPEI, TAIWAN, AND NCTS (TAIPEI OFFICE)

*E-mail address:* chiafu@math.sinica.edu.tw

(J.-D. Yu) DEPARTMENT OF MATHEMATICS, NATIONAL TAIWAN UNIVERSITY, TAIPEI, TAIWAN

*E-mail address:* jdyu@math.ntu.edu.tw